

Mu Alpha Theta National Convention 2003

Calculus Individual Test

For all questions, answer E. "NOTA" means none of the above answers is correct.

1. The total cost, c , to a company for selling x "Barker's Backyard Barbeque Grills" is $c(x) = x^2 + 4x + 100$. The selling price per grill is $p(x) = 100 - x$. What price per grill will yield the maximum profit for the company?

A. 98 B. 76 C. 50 D. 24 E. NOTA
2. For $x > 0$, $\int \frac{2x + x^2 - \sqrt{x}}{x^2} dx =$

A. $\frac{2 + 2\sqrt{x} \ln x + x\sqrt{x}}{\sqrt{x}} + C$ B. $\frac{1 + 4\sqrt{x} \ln x + 2x\sqrt{x}}{2\sqrt{x}} + C$

C. $\frac{2x^2 + 4\sqrt{x} + x^3\sqrt{x}}{x^2\sqrt{x}} + C$ D. $\frac{2 + 2\sqrt{x} \ln x}{\sqrt{x}} + C$ E. NOTA
3. Find the slope of the curve $y = 8 - x - x^2$ at its second quadrant point of intersection with the line $x - 2y + 7 = 0$.

A. -4 B. -2 C. 2 D. 5 E. NOTA
4. The radius of a circle increases from 2.00 cm to 2.02 cm. Use differentials to estimate the resulting change in the circle's area.

A. $\frac{2\pi}{25}$ B. $\frac{4\pi}{25}$ C. 4π D. $\frac{\pi}{25}$ E. NOTA
5. Use the Trapezoidal Rule with $n = 3$ to approximate the area under the curve $y = \cos x$ and above the x -axis from $a = 0$ to $b = \frac{\pi}{2}$.

A. $\frac{\pi(2 + \sqrt{3})}{12}$ B. $\frac{\pi(1 + \sqrt{3})}{12}$ C. $\frac{\pi(1 + \sqrt{3})}{6}$ D. $\frac{\pi(2 + \sqrt{3})}{6}$ E. NOTA
6. Consider the function $g(x) = 3x^2 e^x$. How many of the following statements regarding $g(x)$ is/are true?

I. $g(x)$ is positive for all real values of x .
 II. $g(x)$ is decreasing on the interval $(0, 2)$.
 III. $g(x)$ has exactly one point of inflection.
 IV. $g(x)$ has a relative minimum at $x = -2$.

A. 1 B. 2 C. 3 D. 4 E. NOTA
7. Let $u(x) = \sqrt{x^2 + 9}$ and $v(x) = 3x^3 - 2x$. Find $\frac{du}{dv}$ evaluated at $x = 4$.

A. $\frac{1}{1420}$ B. $\frac{3}{755}$ C. $\frac{4}{855}$ D. $\frac{2}{355}$ E. NOTA

8. Let f be a function such that $f(x+h) - f(x) = 6xh + 3h^2$ and $f(1) = 5$. Determine $f(2) + f'(2)$.
- A. 8 B. 13 C. 16 D. 26 E. NOTA
9. If $f(x) = \ln(2x+3)$ for $x > -\frac{3}{2}$, then the n th derivative of f at x equals
- A. $\frac{-(2)^n (n-1)!}{(2x+3)^n}$ B. $\frac{(-1)^{n+1} (n-1)!}{(2x+3)^n}$ C. $\frac{-(-2)^n (n-1)!}{(2x+3)^n}$ D. $\frac{2^n}{(2x+3)^n}$ E. NOTA
10. An equation of the tangent line to the graph of a differentiable function g at $x=0$ is $y+1=4(x-0)$. Evaluate $\lim_{x \rightarrow 0} \frac{\sin(3x)}{2x g(x)}$.
- A. $-\frac{3}{2}$ B. $\frac{1}{8}$ C. $\frac{3}{8}$ D. $\frac{1}{2}$ E. NOTA
11. The value of a particular investment at time t is increasing at a rate proportional to the difference between \$20,000 and its value at t . At $t=0$, the value is \$2000 while at $t=1$, the value is \$3000. Find the value (in dollars) at $t=3$.
- A. $20,000 - 17,000\left(\frac{17}{18}\right)^2$ B. $20,000 - 18,000\left(\frac{17}{18}\right)^2$
 C. $20,000 - 17,000\left(\frac{17}{18}\right)^3$ D. $20,000 - 18,000\left(\frac{17}{18}\right)^3$ E. NOTA
12. Let $g(x) = \int_2^{3x} \sin^2 t \, dt$. Find $g'''(x)$.
- A. $54 \cos(6x)$ B. $-162 \cos(3x) \sin(3x)$
 C. $-18 \sin^2(3x) + 18 \cos^2(3x)$ D. $-216 \sin(3x) \cos(3x)$ E. NOTA
13. Let s be the length of each one of the congruent sides of an isosceles triangle and let θ be the angle between them. If s is decreasing at the rate of $\frac{1}{10}$ ft/min and θ is increasing at the rate of $\frac{\pi}{90}$ radians/min, then at what rate, in ft^2/min , is the area of the triangle changing when $s = 2\sqrt{3}$ ft and $\theta = \frac{\pi}{6}$?
- A. $\frac{\sqrt{3}\pi}{30} + \frac{\sqrt{3}}{10}$ B. $\frac{1}{60}(\pi + 6\sqrt{3})$ C. $\frac{1}{30}(\pi - 9)$ D. $\frac{\sqrt{3}}{30}(\pi - 3)$ E. NOTA
14. Find $\frac{dy}{dx}$ for the relation $\sin x = x + x \tan y$
- A. $\frac{1 + \tan y - \cos x}{x \sec^2 y}$ B. $\frac{\cos x - 1 - \tan y}{x + x \sec^2 y}$ C. $\frac{x \cos x - \sin x + x}{x^2 \sec^2 y}$ D. $\frac{x \sin x - \cos x}{x^2 + x^2 \sec^2 y}$ E. NOTA

15. The function $a(t) = (t - 8)^{-2/3}$ for $t \geq 0$ represents the acceleration (in ft/sec²) of a moving body whose velocity after 9 seconds is 6 ft/sec. Find the total distance traveled (in feet) by the body during the first 8 seconds.
- A. 12 B. $\frac{27}{2}$ C. 36 D. 288 E. NOTA
16. Given that $f''(x) = 6|x|$, which of the following functions could be $f(x)$?
- I. $|x^3|$ II. $x \cdot |x^2|$ III. $\frac{|x^4|}{x}$
- IV. $|x| \cdot x^2$ V. $\frac{x^4}{|x|}$
- A. I, II & III only B. I & IV only C. I, IV & V only D. II, III & V only E. NOTA
17. The "Mean Value Theorem for Integrals" states that for a continuous function $f(x)$ on the interval $[a,b]$, there is at least one value c in the interval for which $f(c)$ is equal to the average value of the function on the interval. Given the function $f(x) = \begin{cases} x^3 & \text{if } 0 \leq x \leq 1 \\ x^2 & \text{if } 1 < x \leq 3 \end{cases}$ on the interval $[0,3]$, find all values of c guaranteed by the theorem.
- A. $\sqrt[3]{\frac{107}{36}}$ only B. $\sqrt[3]{\frac{107}{12}}$ only C. $\frac{\sqrt{107}}{6}$ only D. $\frac{\sqrt{107}}{2\sqrt{3}}$ only E. NOTA
18. Given $h(x) = x^2$, $g(x) = h(1 + f(x))$ and $f'(1) = g'(1) = 1$. Find $f(1)$.
- A. -1 B. $-\frac{1}{2}$ C. 0 D. 1 E. NOTA
19. Let R_1 be the plane region bounded by the curves $y = \sqrt{x^3}$, $y = 0$ and $x = 4$. Let R_2 be the plane region bounded by the curves $y = \sqrt{x^3}$, $x = 0$ and $y = 8$. Let V_1 be the volume of the solid formed when R_1 is revolved about the x -axis. Let V_2 be the volume of the solid formed when R_2 is revolved about the x -axis. Find the ratio of V_1 to V_2 .
- A. 1 : 2 B. 1 : 3 C. 1 : 4 D. 1 : 6 E. NOTA
20. To which of the following functions does Rolle's Theorem apply?
- A. $f(x) = \frac{x^2 - 1}{x}$ on $[-1,1]$ B. $g(x) = 4 + |x - 2|$ on $[0,4]$
- C. $h(x) = x^2 - 2x$ on $[-2,0]$ D. $k(x) = 4 - \frac{3}{x}$ on $[1,3]$ E. NOTA

21. Evaluate $\int \frac{5x \, dx}{\sqrt[3]{(x^2 + 7)^2}}$

A. $\frac{5}{6} \sqrt[3]{x^2 + 7} + C$

B. $\frac{3}{10} \sqrt[3]{x^2 + 7} + C$

C. $\frac{15}{2} \sqrt[3]{x^2 + 7} + C$

D. $\frac{1}{30} \sqrt[3]{x^2 + 7} + C$

E. NOTA

22. An index of consumer confidence fluctuates between -1 and 1 . Over a two-year period, beginning at time $t = 0$, the level of this index, c , is $c(t) = \frac{t \cos(t^2)}{2}$, where t is measured in years. Find the average value of this index over the two-year period.

A. $-\frac{1}{4} \sin(4)$

B. 0

C. $\frac{1}{4} \sin(4)$

D. $\frac{1}{2} \sin(4)$

E. NOTA

23. Suppose f is continuous and $f(x) = 1 + \int_0^x [(f(t))^2 + 1] \, dt$ for $0 \leq x \leq \frac{1}{2}$. Which of the following functions could be $f(x)$ on $[0, \frac{1}{2}]$?

I. $\frac{1}{x^2 + 1}$

II. $\arctan(x) + 1$

III. $\tan(x) + 1$

IV. $\tan(x + \frac{\pi}{4})$

A. I & IV only

B. II & III only

C. III only

D. IV only

E. NOTA

24. Find the equation of the line joining the inflection points of $y = \frac{2 - x}{x^2 + 4}$.

A. $3x + 16y = 2$

B. $x + 16y = 6$

C. $3x + 8y = 4$

D. $x + 8y = 12$

E. NOTA

25. Evaluate $\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{n}{n^2 + k^2}$

A. $\ln 2$

B. $\frac{\sqrt{3}}{2}$

C. $\frac{\pi}{4}$

D. 1

E. NOTA

26. Let $f(x)$ equal the function whose derivative at any value of x is $f'(x) = -2x e^{-x^2}$ and satisfying the condition $f(0) = 1$.

Let $g(x)$ equal $\lim_{h \rightarrow 0} \frac{\int_1^{x+h} e^{-t^2} dt - \int_1^x e^{-t^2} dt}{h}$.

Let $h(x)$ equal $\frac{d}{dx} \left(\frac{-e^{-x^2}}{2x} \right)$.

Which of these functions are equal?

- A. f and g only B. g and h only C. f and h only D. f , g and h E. NOTA

27. Let $f(x) = \begin{cases} 3x^2 & \text{for } 0 \leq x \leq 1 \\ 4 - x & \text{for } 1 < x \leq 4 \end{cases}$ and let R be the region bounded by the graph of f , the x -axis, and the lines $x = b$ and $x = b + 2$, where $0 \leq b \leq 1$. Determine the value of b that maximizes the area of R .

- A. $\frac{1}{2}$ B. $\frac{2}{3}$ C. $\frac{3}{4}$ D. 1 E. NOTA

28. A disease is spreading through a population in a manner that can be modeled by the function $f(t) = \frac{P}{1 + 3e^{-t}}$ where P is the total population and $f(t)$ is the number infected at time t . What proportion of the population is infected when the disease is spreading the fastest?

- A. $\frac{1}{3}$ B. $\frac{1}{e}$ C. $\frac{1}{2}$ D. $\frac{2}{3}$ E. NOTA

29. Which one of the following is an equation of the line with slope $\frac{25}{2}$ that is normal to the graph of $y = \frac{1}{(5x + 2)^2}$?

- A. $625x - 50y = 373$ B. $625x - 50y = 371$
 C. $625x - 50y = 367$ D. $625x - 50y = 361$ E. NOTA

30. Evaluate $\int_{1/2}^1 3x\sqrt{2x-1} dx$

- A. $\frac{233}{960}$ B. $\frac{4}{15}$ C. $\frac{233}{320}$ D. $\frac{4}{5}$ E. NOTA